

Impact of processing upon dry blueberry fruit aqueous extracts

Silva, S.¹; Costa, E. M. ¹; Costa, M. R.¹; Pereira, M. F. ¹; Morais, M. R. ¹; Pintado, M. E. ¹

¹CBQF/Escola Superior de Biotecnologia, Universidade Católica Portuguesa, Rua Dr. António Bernardino de Almeida, Porto, Portugal




Introduction

The *Vaccinium* genus, from the *Ericaceae* family, is composed by ca. 450 species distributed worldwide. *Vaccinium corymbosum*, also known as highbush blueberry, is one of the several species that are known to produce blueberries (small, dark blue berries with a bittersweet taste). Common cultivated varieties are either Northern Highbush Blueberries (NHB) or Southern Highbush Blueberries (SHB) depending on their temperature requirements. Blueberries are known as a good source of phenolic compounds. Kader *et al.* (1996), Skrede *et al.* (2000) and Brambilla *et al.* (2008) reported three types of phenolic acids (gallic, syringic and vanillic) and four types of cinnamic acids (chlorogenic, the major cinnamic derivative, caffeic, ferulic, o- and p-coumaric). Due to the high content in pectins found in blueberry the standard procedure in industry is to depectinize the blueberry prior to its use as an ingredient. Considering that one of the possible applications of these extracts is as a functional ingredient, the aim of this study was to access the impact of depectinization when performed after processing and pectinases were inactivated through heat treatment.



Procedure



Results




Fig.1 – Total antioxidant capacity.




Fig.2 – Total phenolic content.




Fig.3 – Total anthocyanin content.




Fig.3 – Chromatographic profile at 520 (2) and 310 (1) nm for untreated (a) and depectinized (b) infusions.




Fig.4– Chromatographic profile at 520 (2) and 310 (1) nm for untreated (a) and depectinized (b) boiled fruit extracts.

Conclusions

- Depectinization showed an impact on total antioxidant capacity and total anthocyanin's content of fruit infusions
- Depectinization had no apparent effect upon boiled fruit extracts
- Chromatographic profiling showed a considerable alteration of the anthocyanins profile for both extraction types after depectinization
- Chromatographic profiling showed little variation of the phenolic profile for both extraction types after depectinization

References

- Brambilla, A.; Lo Scalzo, R.; Bertolo, G.; Torreggiani, D. J AGR FOOD CHEM 2008 56(8): 2643-2648
- Kader, F.; Rovel, B.; Girardin, M.; Metche, M. FOOD CHEM 1996 55: 35-40
- Skrede, G.; Wrolstad, R.E.; Durst, R.W. J FOOD SCI 2000 65(2): 357-364



Acknowledgements

The authors hereby acknowledge the Agency of Innovation (Agência de Inovação, ADI, Portugal) and Quadro de Referência Estratégico Nacional (QREN, Portugal) which through the project "Myrtillus – Mirtillo com inovação" (QREN-ADI 13736) provided funding for the realization of this work and to the project PEst-OE/EQB/LA0016/2011.